

# Magnitudes and implications of peak discharges from glacial Lake Missoula

JIM E. O'CONNOR\* }  
VICTOR R. BAKER } Department of Geosciences, University of Arizona, Tucson, Arizona 85721

## ABSTRACT

New field evidence and discharge calculation procedures provide new estimates of maximum late Pleistocene glacial Lake Missoula flood discharges for two important reaches along the flood route. Within the Spokane Valley, near the point of release, the peak discharge probably exceeded  $17 \pm 3$  million  $\text{m}^3 \cdot \text{sec}^{-1}$ . Downstream at Wallula Gap, a major point of flow convergence, peak discharge was about  $10 \pm 2.5$  million  $\text{m}^3 \cdot \text{sec}^{-1}$ . Flow duration was on the order of several days. These are the largest known terrestrial fresh-water flows.

Consideration of these discharge values constrains models for the failure of glacial Lake Missoula. The maximum discharges estimated here are larger than theoretical and empirical predictions of maximum subglacial jökulhlaup-style releases for Lake Missoula. We postulate, consistent with geological relations in the glacial Lake Missoula basin and in the Channeled Scabland, that the largest late Wisconsinan Missoula Flood resulted from a cataclysmic failure of the impounding ice dam of glacial Lake Missoula. This large release may have been the result of a complete rupture of the ice dam. Subsequent multiple flows of lesser magnitude may have resulted from repeated subglacial releases from the lake.

## INTRODUCTION

Pleistocene discharges from ice-dammed glacial Lake Missoula produced a suite of spectacular flood features along multiple flow pathways between western Montana and the Pacific Ocean (Fig. 1). Since the first decades of this century, scientific controversy has continued regarding the interpretation of these features and

deposits, including the origin, chronology, magnitude, frequency, and lake-release mechanisms associated with possible repeated emptying of Lake Missoula. Baker and Bunker (1985) and Waitt (1985) reviewed the development of hypotheses regarding these issues. Here we report results of field studies and hydraulic reconstructions of maximum discharges associated with the Missoula floods at two important sites, and we discuss the implications of these results to hypotheses proposed and models advanced concerning the nature of Lake Missoula flooding.

## METHODS

The primary purpose of this study is to provide limits, based on physical geologic evidence of flood stages, on the maximum discharges that issued from glacial Lake Missoula. Field studies were conducted in the Wallula Gap reach where there is little reported information regarding maximum flood stages. Further studies were also performed in the Rathdrum Prairie and Spokane Valley, where existing information (Baker, 1973) was re-evaluated and augmented. The evidence for maximum stages is combined with step-backwater flow modeling to estimate peak discharges through each of the reaches.

## Geologic Evidence of Maximum Flood Stages

Altitudes of flood landforms and deposits provide minimum values for the maximum flow stages, because the depth of water above the flood feature is unknown. In contrast, nearby nonflooded terrains indicate the maximum possible local altitude of peak flow stage, because the actual water surface must have been lower. Evidence of flooding includes divide crossings, "scabland" topography, scarps cut into loess, ice-rafted erratics, and accumulations of flood-transported sediment (Baker, 1973, p. 13-14). Divide crossings, loess scarps, and scabland topography are most useful as paleostage indicators in reaches of relatively steep flow gradient

and can generally be recognized on topographic maps and aerial photographs. Ice-rafted erratics, exotic rock types beyond the limits of glaciation, are the most consistent indicators of maximum flood stages in ponded areas. Field studies were undertaken in both study reaches to confirm map and photo interpretations, to evaluate existing data, and to determine the narrowest possible altitude range of maximum flow stages. Altitudes of stage evidence were determined by mapping onto U.S. Geological Survey 7.5-minute topographic maps. Altitudes were determined conservatively for features identified in this study and those of Baker (1973); that is, positions inferred to be flooded were assigned altitudes on the basis of the highest contour line determined to be below that position. Therefore, the reported altitude of maximum flow stage is as much as one contour interval (6 to 12 m) below the maximum flow stage. Likewise, reported altitudes of features inferred not to have been inundated are as much as one contour interval above the inferred maximum flow stage.

## Discharge Calculations

**Step-Backwater Modeling.** To estimate discharges associated with the field evidence of maximum stages, we used the U.S. Army Corps of Engineers' HEC-2 Water Surface Profiles computer program (Feldman, 1981; Hydrologic Engineering Center, 1985). This algorithm computes energy-balanced water-surface profiles for steady, gradually varied flow on the basis of the one-dimensional energy equation (a form of the Bernoulli Equation) with the step-backwater method of profile computation (Chow, 1959, p. 265-268). For flow conditions (discharge, stage, and channel geometry) specified at one cross section, stage is calculated for an adjacent cross section of known geometry and reach length by conserving mechanical energy and accounting for estimated flow-energy losses associated with roughness and channel expansions and contractions. Calculated

\*Present address: U.S. Geological Survey, David A. Johnston Cascades Volcano Observatory, 5400 MacArthur Boulevard, Vancouver, Washington 98661.




Figure 1. Major features associated with Missoula flooding in eastern Washington, northern Idaho, and western Montana. Geology from Baker (1973), Waite and Thorson (1983), and Atwater (1986).

conditions at the second cross section are then taken as "known" for the next incremental subreach. In this manner, the stage can be determined for each cross section, and the water-surface profile for a specified discharge can be determined for an indefinite length of channel. For subcritical flow, profiles are calculated successively in the upstream direction.

**Application for Paleohydrologic Analysis.** For geologic applications, where the object is to estimate the discharge associated with paleostage evidence, water-surface profiles are computed for different combinations of discharge values and energy loss coefficients until a calculated profile is achieved that most closely matches the water-surface profile indicated by the maximum stage evidence (Ely and Baker, 1985; Craig and Hanson, 1985; Jarrett and Malde, 1987; O'Connor and Webb, 1988). This method of flow retrodiction probably represents an improvement over most previous discharge

calculations in the Channeled Scabland for the following reasons. (1) It does not require the assumption of uniform flow in a highly nonuniform environment as does simple application of the Chezy Equation (Bretz, 1925). (2) In contrast to the slope-area method (Dalrymple and Benson, 1967; Baker, 1973, p. 17-18), computed step-backwater profiles are not dependent on the precise altitudes of the stage evidence; therefore, discharge estimates are not particularly sensitive to local uncertainties in actual stage altitudes.

Assumptions intrinsic to step-backwater analyses are that the flow is steady, gradually varied, and one-dimensional. The study reaches were selected partly to minimize deviations from these assumptions; the following reach descriptions discuss the suitability of each site for step-backwater calculations. Additional assumptions are introduced for flows at this scale and for flows that occurred in the past. Importantly, it is

uncertain whether using the empirical Manning Equation to evaluate local friction slope (intrinsic to most indirect methods of discharge estimation) is appropriate for flows that are several orders of magnitude larger than any flow that has been directly measured. We know of no way to evaluate this assumption except to note that the Manning Equation has proven to be appropriate for flows over several (albeit much smaller) orders of magnitude. Another assumption is that the existing measured cross sections closely approximate the channel geometry at peak flood stages; that is, there hasn't been significant post-peak downcutting, filling, or valley widening. This assumption must also be evaluated at each site, as described below.

For both reaches, cross sections were measured from U.S. Geological Survey topographic maps and were positioned and spaced (1) to characterize downstream changes in channel geometry and (2) to limit changes in flow condi-

tions between sections. Manning roughness coefficients were assigned for appropriate portions of each cross section according to the estimated degree of flow efficiency across the surface. Although there is no rational method available to determine roughness coefficients for flows of this scale, the chosen values were selected on the basis of systematic criteria and experience with modeling similar flows (O'Connor, 1990). Sensitivity tests were performed to estimate the ef-

fect of uncertainties in roughness coefficients on the resultant water-surface profiles. Flow energy losses resulting from flow expansions and contractions (form roughness) are calculated as being proportional to the change in kinetic energy from one cross section to another (Hydrologic Engineering Center, 1985). Different values of these coefficients were also used to test their effect on the calculated water-surface profiles, as described below.

**STUDY REACHES**

**Wallula Gap**

Wallula Gap is a narrow, 1.5-km-wide, 250-m-deep gorge through the Horse Heaven Hills anticline in south-central Washington (Fig. 2). The constriction is the southern boundary of the Pasco Basin, a basin that has accumulated as much as 300 m of late Tertiary and Quaternary




Figure 2. Topography, locations of evidence for maximum stages (Table 2), cross-section locations, and approximate area of inundation for the Wallula Gap reach at maximum flood stages. Note that some of the evidence for maximum stages is outside the map area.

TABLE 1. DISCHARGE ESTIMATES FOR THE MISSOULA FLOODS

Location	Discharge (10 <sup>6</sup> m <sup>3</sup> · sec <sup>-1</sup> )	Reference	Calculation procedure	Comments
Wallula Gap	1.9	Bretz (1925, p. 257-258)	Chezy equation	Calculation performed by D. F. Higgins. Utilized cross-section areas much smaller than those associated with present understanding of maximum flood stages.
Wallula Gap	9.1	Baker (1973)	Contracted opening equation	Assumed a maximum ponding level of 350 m in the Pasco Basin. Assumed critical flow through Wallula Gap.
Wallula Gap	12.5	Craig and Hanson (1985, p. 52-53)	Step backwater	Assumed a maximum ponding level of 350 m in the Pasco Basin. Assumed critical flow through Wallula Gap.
Wallula Gap	10	This study	Step backwater	Profile constraints include 335-m ponding level in the Umatilla Basin and a 370- to 385-m ponding level in the Pasco Basin. Flow not assumed to be critical in the constriction.
Sentinel Gap	9.5	Craig and Hanson (1985, p. 53-66)	Step backwater	Downstream stage estimated on the basis of an extrapolated water-surface slope from Wallula Gap. Upstream ponding level assumed to be 366 m. Discharge estimate applies only to water that followed the course of the Columbia River (does not include flow in the eastern scabland tracts).
Moses Coulee	1.1-9.8	Hanson (1970, p. 59)	Manning equation	Limiting values considering estimated maximum and minimum cross-section areas at peak stages for two different cross sections.
Grand Coulee (near Soap Lake)	4.5	Baker (1973, Pl. 1)	Contracted opening equation	Assumed critical flow through the constriction at Soap Lake. Upstream water surface estimated from geologic evidence of maximum stage.
Grand Coulee (near Steamboat Rock)	0.13	Atwater (1986)	Sediment transport competence criteria	Based on extrapolation of a local velocity estimate across the upper Grand Coulee. Discharge associated with flood deposits within glacial Lake Columbia sediments.
Cheney-Palouse scabland tract	8.1	Baker (1973)	Slope-area method	Based on water-surface slope defined by geologic evidence of maximum stages.
Spokane Valley-Rathdrum Prairie	21.3	Baker (1973)	Slope-area method	Based on water-surface slope defined by geologic evidence of maximum stages.
Spokane Valley-Rathdrum Prairie	17	This study	Step backwater	Based on highwater evidence of Baker (1973) and additional field observations.
Lake Missoula breakout point	13-15.3 2.6-3.5	Clarke and others (1984)	Simulation model	Model simulating subglacial release of Lake Missoula from the maximum lake level. High values are for conditions of no downstream backwater and immediate conveyance of lake water to the release point. Low values are for conditions of downstream backwater and delayed conveyance associated with the complex lake geometry.
Lake Missoula breakout point	2.1	Beget (1986)	Empirical relationship	Extrapolated relationship between ice-dammed lake volumes and associated jökulhlaup discharges to predict discharge associated with the release of the maximum volume of Lake Missoula.
Eddy Narrows	10.9	Pardee (1942)	Chezy and Manning equations	Used high-level flood features to estimate cross-section area. Used present channel slope as a proxy for the water surface (and energy slope).

fluvial and lacustrine sediments (Newcomb, 1958; Newcomb and others, 1972; Waitt and Swanson, 1987), including glacial Lake Missoula flood deposits. Wallula Gap is an important site for discharge estimation, because it is the most upstream point of convergence for the multitude of flow routes that enter the Pasco Basin (Fig. 1). Previous discharge estimates at Wallula Gap were calculated by Bretz (1925), Baker (1973), and Craig and Hanson (1985) (Table 1).

Wallula Gap is particularly suitable for flow reconstruction by the step-backwater method. The steady-flow assumption is probably satisfied because of hydraulic damming at the constriction and consequent upstream storage and flow attenuation within the Pasco Basin. A radiometrically dated basalt flow close to present river level near The Dalles indicates that the Columbia River has been near its present altitude for at least 900,000 yr (Bjornstad and others, 1991), and flood-related downcutting was minor. Because the constriction is formed in resistant rocks, flood-related widening probably does not affect discharge estimates. As noted by Bretz (1969), the existence at river level of channel-margin flood sediments associated with maximum stages indicates that there has not been major channel widening after the passing of maximum discharges. Bretz (1969, p. 535) further stated, "[The] gorge [immediately] downstream from the gap itself therefore was deep as it is today. . . ."

Previous workers in the area have defined, on the basis of ice-rafterd erratics, maximum flood stages in the Pasco Basin (370-385 m; Bjornstad and others, 1991) and in the Umatilla Basin at the downstream end of the study reach (335 m;




TABLE 2. MAXIMUM STAGE EVIDENCE FOR THE WALLULA GAP REACH

Number	Location (and USGS 7.5' quadrangle)	Altitude (m)	Type	Reference	Comments
1	W½ S26 T. 2 N. R. 25 E. Strawberry Canyon SW (OR)	320	Ice-rafted erratic	Allison (1933, p. 689)	Altitude as reported by Allison.
2	Umatilla Basin	335-350	Fine-grained deposits and small erratics	Allison (1933, p. 685)	Upper limit of "thin cover of the pebbly and bouldery glacial silt." Lower altitude estimate indicated in Figure 4. (We have found such evidence of flooding only to an altitude of 335 m.)
3	SW¼ SW¼ S17 T. 6 N. R. 28 E. Umatilla (WA-OR)	305	Fluvial sands and silts	This study	Felsic boulders (ice-rafted?) at similar altitudes on adjacent hillslopes.
4	NW¼ S8 T. 2 N. R. 30 E. Nolin (OR)	310	Ice-rafted erratic	Allison (1933, p. 685)	Altitude as reported by Allison.
5	NW¼ NE¼ S17 T. 2 N. R. 30 E. Nolin (OR)	300	Ice-rafted erratics	This study	
6	W½ S13 T. 5 N. R. 30 E. Juniper (OR-WA)	323	Scabland topography	This study	Stripped basalt and eroded loess.
7	NE¼ NE¼ S28 T. 6 N. R. 30 E. Juniper (OR-WA)	315	Base of loess scarp	This study	Top of loess scarp is at 342 m and provides upper limit for maximum flood stage.
8	SW¼ SW¼ S30 T. 6 N. R. 31 E. Juniper Canyon (OR-WA)	349	Top of loess scarp (upper limit)	This study	Non-flooded loess-covered terrain.
9	NW¼ SW¼ S30 T. 6 N. R. 31 E. Juniper Canyon (OR-WA)	305	Divide crossing	This study	
10	NW¼ NE¼ S13 T. 6 N. R. 30 E. Nine Canyon (WA)	329	Divide crossing	This study	
11	SW¼ SW¼ S6 T. 6 N. R. 30 E. Wallula (WA-OR)	366	Divide not crossed (upper limit)	This study	
12	NE¼ SE¼ S6 T. 6 N. R. 31 E. Wallula (WA-OR)	343	Un-eroded loess-covered mesa (upper limit)	This study	
13	SW¼ NE¼ S5 T. 6 N. R. 31 E. Wallula (WA-OR)	336-349	Scabland topography	This study	Boulder deposits to an altitude of 336 m.
14	NW¼ NW¼ S5 T. 7 N. R. 31 E. Wallula (WA-OR)	360	Un-eroded loess-covered mesa (upper limit)	This study	
15	SW¼ SW¼ S32 T. 7 N. R. 31 E. Wallula (WA-OR)	342	Divide crossing	This study	
16	SW¼ NE¼ S32 T. 7 N. R. 31 E. Wallula (WA-OR)	342	Ice-rafted erratics	This study	
17	SW¼ SW¼ S35 T. 7 N. R. 31 E. Wallula (WA-OR)	355	Boulder bar and divide crossing	This study	Altitude is of boulder bar.
18	SW¼ SW¼ S35 T. 7 N. R. 31 E. Wallula (WA-OR)	378	Top of loess scarp (upper limit)	This study	Immediately above 17.
19	SW¼ SW¼ S30 T. 7 N. R. 31 E. Wallula (WA-OR)	348-366	Divide crossing	Bretz (1969); this study	Bottom of divide is at 348 m. Bretz postulated a maximum flood stage of 366 m.
20	Pasco Basin	370-385	Ice-rafted erratics	Bjornstad and others (1991)	Maximum ponding level in the Pasco Basin.

Allison, 1933). These values are consistent with our observations of maximum stages inferred from the field evidence within Wallula Gap (Table 2).

**Wallula Gap Modeling and Results.** Flow profiles were calculated through 21 cross sections over a length of 30 km (Figs. 2 and 3). The maximum flow stage was assumed to be 335 m at the downstream cross section. Manning roughness coefficients were assigned as follows: 0.04 over the valley floor; 0.05 over the steep valley margins; 0.1 for flow over the high uplands. These values are larger than roughness coefficients typically assigned to more normal scale floods. Following Jarrett and Malde (1987), in their application of step-backwater modeling techniques to the Bonneville Flood, we used contraction-expansion coefficients of 0.0 and 0.5. With these parameters, a discharge of 10 million  $\text{m}^3\text{sec}^{-1}$  provides the best fit, by eye, to the geologic evidence of maximum Missoula Flood stages through Wallula Gap and results in a ponding level of 375 m in the Pasco Basin (Fig. 3). Utilizing the same energy-loss coefficients, an 11.2 million  $\text{m}^3\text{sec}^{-1}$  discharge is required to achieve the 385-m ponding elevation reported by Bjornstad and others (1991) as an upper estimate of maximum ponding levels in the Pasco Basin. In our estimation, the 10 million  $\text{m}^3\text{sec}^{-1}$  estimate provides a better fit to the maximum stage evidence through the constriction and serves as a minimum estimate for the energy-loss coefficients specified above. For comparison, the maximum measured discharge for the Columbia River was 35,000  $\text{m}^3\text{sec}^{-1}$  at The Dalles during the 1894 flood.

The 10 million  $\text{m}^3\text{sec}^{-1}$  discharge estimate is minimally affected by uncertainties in the Manning roughness coefficients. Varying Manning roughness coefficients by  $\pm 25\%$  requires adjustments of less than 5% to the prescribed discharge value to maintain a 375-m stage in Pasco Basin (Fig. 4). Modeling results are more sensitive to the expansion and contraction coefficients. Increasing these coefficients to the values of 0.3 and 0.7, values larger than typically assigned for estimating this form of energy loss, the required discharge to achieve a 375-m ponding level decreases 20% to about 8 million  $\text{m}^3\text{sec}^{-1}$ . Considering uncertainties in the maximum stages achieved in the Pasco Basin and uncertainties in assigning energy-loss coefficients, discharge values between 7.5 and 12 million  $\text{m}^3\text{sec}^{-1}$  are consistent with maximum stage evidence at the upstream and downstream ends of the reaches (Fig. 4).

Figure 3. Step-backwater calculated water-surface profiles for the Wallula Gap reach. A 10 million  $\text{m}^3\text{sec}^{-1}$  discharge results in a calculated profile that matches most of the highwater evidence along the reach and results in a ponding level in the Pasco Basin consistent with the altitudes of the highest ice-rafted erratics.




Figure 4. Rating curves for calculated stages in the Pasco Basin (cross section 21 of Fig. 3) for different values of energy-loss coefficients and for a ponding level of 335 m in the Umatilla Basin. The effects of varying Manning roughness coefficients are minimal. The principal case was calculated with contraction and expansion energy-loss coefficients of 0.0 and 0.5. Using larger coefficients reduces the discharge required to achieve a 375-m stage in the Pasco Basin to about 8 million  $\text{m}^3\text{sec}^{-1}$ . Considering the highwater evidence identified in this study and the uncertainty of maximum Pasco Basin stages and reasonable ranges of the energy-loss coefficients, the maximum Missoula Flood discharge through Wallula Gap was between 7.5 and 12 million  $\text{m}^3\text{sec}^{-1}$ .

The 10 million  $\text{m}^3\text{sec}^{-1}$  discharge proposed here is somewhat higher than the discharge estimates calculated in the studies by Baker (1973, p. 17–22) and Bretz (1925, p. 257–258) but lower than the 12.5 million  $\text{m}^3\text{sec}^{-1}$  estimate of Craig and Hanson (1985, p. 52–53) (Table 1) (see also Craig, 1987, p. 324). Previous discharge calculations were postulated on assumed ponding levels in the Pasco Basin lower than the maximum level of 370–385 m presently recognized (Bjornstad and others, 1991). Furthermore, the analyses of Baker (1973) and Craig and Hanson (1985) assumed critical-flow conditions within the Wallula Gap constriction. Even though it is likely that critical-flow conditions were achieved during the waxing phase of the flood, it seems that during maximum stages, ponding in the Umatilla Basin to an altitude of 335 m impeded flow through the Wallula Gap. On the basis of the altitudes of a variety of flood features described by Allison (1933) and Newcomb (1969), it appears that constrictions 250 km downstream in the Columbia River Gorge (Fig. 1) hydraulically impounded flow to the

extent that Wallula Gap did not act as a free constriction. If, however, the flood wave traveled at such a speed that downstream backwater effects did not choke the constriction at Wallula Gap until after maximum stages were achieved, then the assumption of critical flow (Baker, 1973; Craig and Hanson, 1985) may be valid. For critical flow conditions within the constriction and a ponding level of 375 m in the Pasco Basin, a discharge of 14 million  $\text{m}^3\text{sec}^{-1}$  could have funneled through Wallula Gap with the energy-loss coefficients specified above (Fig. 5). Because critical flow at Wallula Gap is the most efficient flow condition, this discharge estimate serves as a maximum value, given our present understanding of maximum ponding levels in the Pasco Basin.

#### Spokane Valley and Rathdrum Prairie

The Spokane Valley and the contiguous Rathdrum Prairie is a large, steep-walled valley extending southwest of the southern end of Pend Oreille Lake, Idaho, to Spokane, Washington

(Fig. 6). In the most constricted location (at cross section 5 of Fig. 6), the valley narrows to a width of 6 km before opening up into the Spokane Valley. As much as 150 m of Pleistocene glaciofluvial and flood deposits underlie the valley floor (Bolke and Vaccaro, 1981). Some of these sediments have been molded into flood forms, including bars and trains of "giant current ripples" (Baker, 1973, p. 48–64). Eddy bars, blocking pre-existing valleys, impound many of the lakes marginal to the study reach (Kiver and Stradling, 1982, 1989).

This reach is adjacent to the area of release for the Missoula floods near the south end of Pend Oreille Lake, and the valley served as the primary conduit of flood waters exiting Lake Missoula. As such, discharges within this reach were probably the largest with respect to the entire Channeled Scabland complex. Previous Missoula flood discharge estimates pertinent to this reach include the slope-area calculation of Baker (1973, p. 14–22) based on field evidence of maximum stages, and the discharge calculations for the maximum Lake Missoula release by




Figure 5. A calculated stage-discharge rating curve for the Pasco Basin associated with critical flow conditions within Wallula Gap and the relationship of this curve to the maximum reported altitudes of ice-rafted erratics and rhythmite deposits in the Pasco Basin. The rating curve was calculated by assuming critical flow at cross section 16 (Fig. 3) and computing the upstream profiles with the same (upstream) cross sections used for calculating the water-surface profiles for the entire reach. Assigned energy-loss coefficients were the same as for the principal case of Figure 4. This rating curve should closely approximate maximum possible discharge values associated with the corresponding stages in the Pasco Basin. Craig (1987) provided a similar rating curve based on a different set of cross sections (Craig and Hanson, 1985).






Figure 6. Topography, locations of maximum stage evidence (Table 3), cross-section locations, and approximate area of inundation for the Spokane Valley reach. Note that some of the evidence for maximum stages is outside the map area.

Clarke and others (1984) and Beget (1986) concluded from theoretical and empirical analyses, respectively, of flood releases from ice-dammed lakes (Table 1).

**Evidence of Maximum Flood Stages.** Baker (1973, p. 66-67) reported several sites of maximum stage evidence along this reach, of which several were first observed by P. L. Weis (personal commun. with V. R. Baker, 1969). Many of these were revisited for this study, and their altitudes were re-evaluated. Richmond and others (1965) reported evidence of maximum flood stages along this reach, as did Kiver and Stradling (1989). Additional observations from our field studies have also been incorporated into the analysis (Table 3; Fig. 7). Altitudes of maximum stage evidence at the downstream end of the reach approach 750 m, which is similar to the altitude determined by Atwater (1986, p. 7; 1987) for the maximum flood-swollen stage of glacial Lake Columbia. At the upstream end of the study reach, gravel bars and ice-rafted erratics achieve altitudes near 820 m.

**Discharge Calculation.** The ability to produce reliable discharge estimates here is tempered by the complex flow conditions that must have existed through this reach. Initial flow was assuredly unsteady, perhaps manifested as a dynamic wave (Craig, 1987). Unsteadiness in the flow can lead to discharge overestimation in that the flood features related to the calculated steady-flow water-surface profiles may have been emplaced by the crest of a transient wave moving through the study reach. Many of the high flood features are large constructional landforms, however, that we infer to have required formation times longer than that associated with passage of a dynamic flood wave. Key such features include the eddy bar southeast of Spokane at 726 m (stage indicator no. 3 on Table 3) near the downstream end of the study reach and, at the upstream end, the large bars near Spirit Lake and Round Mountain (stage indicators no. 18 and no. 19) at 799 m. Furthermore, the tortuous geometry of sub-basins and constrictions within glacial Lake Missoula would have regulated outflow from the lake into this reach (Pardee, 1942; Clarke and others, 1984; Craig, 1987) after the initial floodwave had passed. We propose that the well-developed flood features preserved along the study reach represent flow conditions that approached steady. Kiver and Stradling (1989) reported evidence of flood erosion much higher than the stage indicators used in this study. Perhaps this high evidence records the transient effects of a dynamic wave.

Another complicating factor in determining the total glacial Lake Missoula discharge in this reach is that multiple flow routes probably operated at peak stages; at least portions of the latest

TABLE 3. MAXIMUM STAGE EVIDENCE FOR THE SPOKANE VALLEY-RATHDRUM PRAIRIE REACH

Number	Location (and USGS 7.5' quadrangle)	Altitude (m)	Type	Reference	Comments
1	Sanpoil River Valley	750	Ice-rafted erratics	Atwater (1986 p. 7)	Maximum, "flood-swollen" level of Glacial Lake Columbia.
2	W½ S28 T. 26 N. R. 44 E. Spokane NE (WA)	735	Divide crossing	This study	Flood deposits immediately downstream at 730 m.
3	NW¼ SW¼ S30 T. 25 N. R. 44 E. Spokane NE (WA)	726	Eddy bar	This study	
4	NE¼ SE¼ S30 T. 25 N. R. 44 E. Spokane NE (WA)	756	Scabland topography	This study	
5	W½ SW¼ S10 T. 24 N. R. 44 E. Freeman (WA)	743	Divide crossing	Baker (1973, no. 61); this study	
6	SW¼ NE¼ S12 T. 24 N. R. 44 E. Freeman (WA)	744	Ice-rafted erratic	This study	
7	SE¼ S27 T. 25 N. R. 45 E. Liberty Lake (WA)	762-775	Divide crossing	Baker (1973, no. 63); this study	Bottom of the divide at 762 m. Maximum altitude of rounded boulders is 775 m.
8	SE¼ SE¼ S24 T. 25 N. R. 45 E. Liberty Lake (WA-ID)	781	Divide crossing	Baker (1973, no. 64); this study	Small gravel delta downstream of divide.
9	SW¼ SW¼ S16 T. 26 N. R. 45 E. Greenacres (WA)	762	Divide crossing	This study	
10	NW¼ SE¼ S21 T. 26 N. R. 45 E. Liberty Lake (WA-ID)	811	Divide not crossed (upper limit)	This study	
11	NE¼ NW¼ S19 T. 50 N. R. 6 W. Liberty Lake (WA-ID)	830	Divide not crossed (upper limit)	Baker (1973, no. 65); this study	
12	NE¼ S36 T. 27 N. R. 45 E. Newman Lake (WA-ID)	799	Divide crossing	This study	
13	SW¼ SE¼ S8 T. 50 N. R. 4 W. Coeur d'Alene (ID)	778	Scabland topography	This study	
14	Coeur d'Alene Lake Basin	813	Ice-rafted erratics	Richmond and others (1965)	Ice-rafted erratics located on "adjacent mountain slopes" flanking Coeur d'Alene Lake.
15	E½ S32 T. 49 N. R. 6 W. Setters (ID-WA)	787	Divide crossing	Richmond and others (1965)	Flow depth unknown.
16	NE¼ NW¼ S32 T. 52 N. R. 3 W. Hayden Lake (ID)	799	Divide crossing	This study	
17	SW¼ NE¼ S17 T. 52 N. R. 3 W. Hayden Lake (ID)	793	Scabland topography	Baker (1973, no. 71)	
18	N½ S29 T. 53 N. R. 4 W. Spirit Lake E. (ID)	799	Flood bar	This study	Large bar with boulders larger than 2 m.
19	SW¼ SW¼ S35 T. 53 N. R. 4 W. Spirit Lake E. (ID)	817	Scabland topography	This study	Upper extent of eroded topography in the vicinity of Round Mountain. Bar downstream is at an altitude greater than 793 m.
20	SE¼ NE¼ S30 T. 53 N. R. 2 W. Bayview (ID)	824	Gravel bar	This study	

Pleistocene floods bypassed the Spokane Valley and Rathdrum Prairie and flowed northward and then south into the Little Spokane River drainage and Spokane basin (Kiver and Stradling, 1982; Waitt, 1984). Flow also bypassed the Spokane Valley by overtopping the Coeur d'Alene Lake basin (Richmond and others, 1965) and spilling southward over divide crossings south of Spokane. Furthermore, during

peak stages, unconsolidated sediment along the valley bottom was probably excavated, creating larger cross-section areas than now present. Water depths approaching 65 m in Coeur d'Alene and Hayden Lakes, lakes that occupy valleys blocked by flood deposits, may provide an indication of the altitude of Rathdrum Prairie when the deposits were emplaced. Flow at maximum stages may have entered Glacial Lake Co-






Figure 7. Calculated profiles for the Spokane Valley reach. A 17 million  $m^3 \cdot sec^{-1}$  is the minimum discharge that results in a calculated profile that matches or exceeds all of the evidence for maximum flood stages. These profiles were calculated on the basis of the present valley topography.

lumbia, a lake formed by the Okanogan ice lobe blocking the Columbia River. At its maximum, stable level of 715 m (Atwater, 1986, p. 6; 1987), ponding extended up the Spokane Valley and inundated most of the study reach.

Because of these complexities and uncertainties, we have focused on determining a *minimum* estimate for the maximum Lake Missoula discharge in this reach. Because we are using the present topography for the flow modeling, calculated cross-section areas are probably smaller than those extant at peak discharge. The high-water indicators used for the flow reconstructions are consistent with the downstream presence of glacial Lake Columbia; however, equally large discharges would have coursed through the study reach at lower stages (at the downstream end of the reach) if driven by a steeper energy gradient associated with unimpounded conditions. The energy-loss coefficients used to calculate the water surface profiles for this reach are larger than those typically used in indirect discharge estimates. As a result, the water surface profile that best matches the stage evidence should be associated with a minimum discharge estimate.


Figure 8. Stage-discharge curves for cross section 12 of the Spokane Valley reach for present topographic conditions. Considering uncertainties in maximum stages and reasonable ranges of energy-loss coefficients, a bracketing range for a minimum estimate of the maximum discharge within this reach is between 14 and 20 million  $m^3 \cdot sec^{-1}$ . For calculations using low energy-loss coefficients, flow conditions were predicted to be near critical at cross section 12.

Water-surface profiles were calculated through 12 cross sections over a distance of 45 km, assuming an initial downstream stage of 750 m. Manning roughness coefficients of 0.04 and 0.1 were chosen to estimate roughness losses over the valley floor and valley walls, respectively. We used contraction/expansion loss coefficients of 0.3 and 0.7.

**Spokane Valley Modeling Results.** For the above parameters and the present topography, a discharge of 17 million  $m^3 \cdot sec^{-1}$  provides the best eye fit to the geologic evidence of maximum stages (Fig. 7). Because the constriction at cross section 5 exerts strong control on the upstream profile at high stages, the presence or absence of glacial Lake Columbia (at a stage of 715 m) has no effect on the discharge value required to em-

place the maximum stage evidence at the upstream end of the reach.

Model variable sensitivity analyses and limits on a minimum estimate for the maximum discharge were postulated on a maximum flow stage of between 817 and 824 m at the upstream end of the reach (constraints imposed by stage indicators 19 and 20), with 824 m assigned as the most likely value of the maximum stage (Fig. 8). Similar to results at Wallula Gap, varying the assigned Manning roughness coefficients by  $\pm 25\%$  has minimal (less than 10%) effect of the calculated discharge. Using smaller values of the expansion and contraction coefficients has a larger effect, raising the discharge required to achieve an 824-m altitude at the upstream end of the reach to about 20 million  $m^3 \cdot sec^{-1}$ . Con-



sidering these ranges of model parameters and uncertainties in the maximum upstream stage, a minimum of 14 to 20 million  $\text{m}^3\cdot\text{sec}^{-1}$  flowed through the Spokane Valley. We emphasize that, because of our approach to reconstructing flow conditions for this reach, this is a *minimum* estimate for *maximum* flow through this reach, as well as being a minimum estimate for maximum glacial Lake Missoula outburst discharges.

This point is substantiated by evaluating the importance of uncertainty in the valley floor altitude at maximum stage. We performed a model trial assuming that the entire valley bottom was excavated to a depth of 60 m relative to its present altitude (consistent with the depths of the marginal lakes). For these conditions, the discharge that produces the required stages is about 27 million  $\text{m}^3\cdot\text{sec}^{-1}$ . Clearly, the valley topography at the time of peak discharge is a major factor in establishing accurate discharge estimates in this reach. It is unlikely, however, that the valley floor was higher than its present altitude during maximum flood stages; hence our 17 million  $\text{m}^3\cdot\text{sec}^{-1}$  estimate is probably a valid minimum estimate.

The 17 million  $\text{m}^3\cdot\text{sec}^{-1}$  discharge determined here is less than the 21 million  $\text{m}^3\cdot\text{sec}^{-1}$  estimate calculated by Baker (1973) with the slope-area method. This discrepancy is probably generated by higher energy-loss coefficients used in this study and the more complete characterization of flow conditions afforded by the step-backwater calculation procedure.

## GENERAL IMPLICATIONS

### Flood Hydrographs

Peak discharge estimates at Wallula Gap and in the Spokane Valley, in conjunction with known storage volumes in Lake Missoula and the Pasco Basin, permit some simple conclusions regarding the flood hydrographs at both localities. For this analysis, we make the following simplifying assumptions. (1) The flood hydrographs were triangular. This is consistent with the general shapes of flood hydrographs produced by dam failures and by some types of jökulhlaups (Costa, 1988). (2) Maximum discharges at both Wallula Gap and in the Spokane Valley were associated with complete emptying of Lake Missoula from its maximum highstand (1,265 m). (3) An insignificant quantity of water bypassed the Spokane Valley flow route. Important physical parameters include the volume of Lake Missoula available for flooding: 2,184  $\text{km}^3$ , according to Clarke and others (1984);




Figure 9. Postulated hydrographs for the largest Missoula Flood(s) (assuming complete emptying of Glacial Lake Missoula from its maximum level). All of the hydrographs represent the same total volume (2,184  $\text{km}^3$ ). a and d are those postulated in this study on the basis of maximum discharges near the outlet and at Wallula Gap. b and c are those proposed by Clarke and others (1984) as resulting from a jökulhlaup release. b represents the extreme case of no "tailwater ponding" (flow impeded by downstream ponding in the Spokane Valley) and immediate conveyance of all of the lake water to the breakout location. c is the more realistic situation of downstream hydraulic conditions affecting the rate of release as well as the delayed response at the breakout point because of the complex lake geometry. e is the hydrograph proposed for Wallula Gap by Craig (1987).

and the volume of hydraulically ponded water behind Wallula Gap: 1,210  $\text{km}^3$  at the 375-m ponding level as determined from the stage-storage curve of Craig (1987).

Constructing a hydrograph for the Spokane Valley reach that satisfies our peak discharge estimate and the volume of Lake Missoula requires a flow duration of about 70 hr (Fig. 9). Because the 17 million  $\text{m}^3\cdot\text{sec}^{-1}$  estimate is a minimum discharge value, it is possible that the actual flow duration was even less. A similar hydrograph can be drawn for Wallula Gap with the added limitation that at peak stage (375 m in the Pasco Basin) there is 1,210  $\text{km}^3$  of ponded storage upstream of Wallula Gap. This imposes a degree of asymmetry to the hydrograph: the waxing part of the hydrograph must have been less than 54 hr, and the waning period was at least 67 hr, with a total flow duration of at least 121 hr (Fig. 9).

Although these hydrographs must be viewed

as first-order approximations, some conclusions can still be drawn. Flow duration associated with such a large discharge was probably remarkably short, on the order of three days within the Spokane Valley. Moreover, the peak discharge seems to have attenuated significantly between the outburst area and Wallula Gap. A large part of peak discharge diminution was probably the result of storage in the Pasco Basin. Stages in the Pasco Basin apparently rose at least 20% faster than they declined. Craig (1987) addressed some of the dynamic aspects of Missoula Flood hydrographs for portions of the flood route, including Wallula gap, and presented hydrographs with shapes that accord with his dynamic model (Fig. 9). Obtaining a complete dynamic picture of the maximum flood(s), however, remains difficult because of the plexus of flow paths across the Channeled Scabland and uncertainty regarding the degree of blockage of the Columbia River valley in north-central

Washington by the Okanogan lobe of the Cordilleran ice sheet.

### The Mode of Ice-Dam Failure

Consideration of the discharge estimates obtained in this study for the Spokane Valley reach may have implications regarding the release mechanism associated with drainage from the highest level of glacial Lake Missoula. In an independent analysis that focused on the dynamics of the failure of the ice dam, Clarke and others (1984) used a physically based computer-simulation model (Clarke, 1982) to predict deductively the range of potential discharges associated with an assumed subglacial style of lake emptying (jökulhlaup) for the glacial Lake Missoula ice dam. Postulating various geometric models and combinations of parameters, they predicted a physically plausible range of maximum discharge values of between 2.5 and 15.3 million  $\text{m}^3\text{-sec}^{-1}$ . For their modeled conditions that most closely approximate the conditions of our step-backwater calculations—downstream stages of 750 m and a more steady contribution of outflow from the entire lake (their conditions of tailwater ponding and a trapezoidal duct conveying outflow to the release area)—Clarke and others (1984) estimated peak outflow rates of 2.6 to 3.5 million  $\text{m}^3\text{-sec}^{-1}$ .

Beget (1986), in commenting on the analysis of Clarke and others (1984), provided an empirical relationship between lake volume and peak discharge based on data from historic jökulhlaups. Extrapolating this relationship to the immensely larger lake volume associated with glacial Lake Missoula, he predicted a maximum Lake Missoula peak discharge of 2.1 million  $\text{m}^3\text{-sec}^{-1}$ . Beget (1986, p. 138) concluded that this estimate “compares favorably with the most conservative and realistic computer model runs of Clarke et al.” and further implied that the field-based discharge estimate of Baker (1973, 21 million  $\text{m}^3\text{-sec}^{-1}$ ) may be an order of magnitude too large.

Implicit in the analyses of Clarke and others (1984) and Beget (1986) is that the ice dam impounding glacial Lake Missoula, at its highest level, failed by buoyant lifting and enlargement of subglacial tunnels. Our best estimate of 17 million  $\text{m}^3\text{-sec}^{-1}$  for the maximum glacial Lake Missoula discharge is a minimum estimate; nevertheless, it is larger than the estimated discharge values associated with a subglacial release, especially the more realistic models. We propose that at least one outburst may have been the result of a more cataclysmic failure of

the ice dam than an outburst resulting from buoyant lifting and subglacial draining. A cataclysmic failure leading to the discharges estimated in this report may have involved complete rupture and removal of the blocking lobe of ice. Although there is no direct evidence of the mode of ice-dam failure, geologic evidence in the Channeled Scabland and in the glacial Lake Missoula basin is consistent with a late Wisconsinan cataclysmic failure of the ice dam followed by a phase of multiple floods associated with periodic subglacial releases.

**Evidence in the Channeled Scabland.** Waitt (1980) originally proposed that a minimum of about 40 late Wisconsinan floods, recorded by sets of rhythmite deposits in the Pasco Basin (commonly referred to in the literature as “Touchet beds”), were jökulhlaups from a regularly self-dumping glacial Lake Missoula. This hypothesis has been found to be consistent with recent interpretations from additional sites of similar stratigraphy preserved in backwater areas along the flood route (Waitt, 1984, 1985). A particularly extensive record is preserved in the Sanpoil River valley, where deposits interpreted to represent at least 89 floods are intercalated within lacustrine deposits of glacial Lake Columbia during blockage of the Columbia River by the Okanogan lobe of the Cordilleran ice sheet (Atwater, 1984, 1986). Interpretation of pertinent radiocarbon dates and varve counts indicates that these floods occurred between  $15,550 \pm 450$  and  $13,050 \pm 650$  yr B.P. Varve counts, flood-bed thickness, and grain-size variations exhibit tendencies consistent with a general up-section decrease in flood magnitudes and recurrence intervals (Atwater, 1986) in a manner consistent with repeated subglacial releases from a thinning ice dam impounding glacial Lake Missoula (Waitt, 1985; Atwater, 1986).

It is apparent from the sedimentologic record that these floods ranged widely in magnitude. Nevertheless, it is not clear that any of the floods recorded in the rhythmite sections is the one that emplaced the highwater evidence used to calculate peak discharge associated with the largest release. For example, in backflooded areas marginal to the Pasco Basin, multiple rhythmites are discernible only to an altitude of 255–260 m (Waitt, 1980, p. 672; 1985, p. 1285; Bunker, 1980, p. 60), whereas bedded silt, interpreted to have been flood-transported but not rhythmically bedded, is found up to altitudes of 350 m (Bretz, 1930, p. 408–415; Flint, 1938, p. 493–494; Newcomb and others, 1972, p. 17, Pl. 1). If a stage of ~260 m was the maximum achieved by the series of flows that deposited the highest

rhythmites, a maximum corresponding discharge value can be estimated from the step-backwater modeling at Wallula Gap. For critical flow conditions in the Wallula Gap constriction, a 5.5 million  $\text{m}^3\text{-sec}^{-1}$  discharge results in a 260-m ponding level in the Pasco Basin (Fig. 5). The corresponding volume of storage in the Pasco Basin is 340  $\text{km}^3$  of water (from the stage-storage relationship of Craig, 1987), less than 30% of the volume associated with a 375-m ponding level.

Pertinent conclusions that can be drawn from the relations in the Pasco Basin are (1) at least one late Wisconsinan flood achieved a stage of 370–385 m, and (2) interpreting each of the highest rhythmites as representing individual floods requires several flows exceeding stages of 255 m. The lack of evidence for multiple flows above 260 m does not rule out higher stages having been achieved repeatedly; however, the disparity between the discharge required to emplace the highest evidence of flooding and those required to emplace the evidence for multiple, repeated flows is consistent with a large flood resulting from a more rapid or complete failure of the ice dam followed by multiple floods that were released by buoyant lifting of the ice dam. A candidate for such a flood may be the large (about 350 m deep near Wenatchee), Wisconsinan-age flood that coursed down the Columbia River prior to complete blockage by the Okanogan lobe of the Cordilleran ice sheet (Waitt, 1977, 1982, 1987). This flood may predate the record of flood sediments in glacial Lake Columbia.

**Evidence in the Glacial Lake Missoula Basin.** Geologic relations in the glacial Lake Missoula basin are also consistent with the idea that there was a single large flood resulting from complete or rapid failure of the ice dam followed by multiple smaller flows that drained subglacially. Pardee (1942) documented cataclysmic flood features associated with a single episode of rapid drawdown from the maximum 1,265- to 1,280-m level of Lake Missoula. Superimposed on some of these features are shorelines indicating refilling of the basin to a maximum altitude of 1,080 m. Pardee (1942, p. 1598) stated, “in this new body of water no unusual currents seem to have developed.” Pardee (1942) ruled out cataclysmic emptying of the younger lake, because of the unmodified nature of the high flood features upon which the shorelines are developed. Chambers (1984) concluded from stratigraphic studies of 40 cycles of bottom sediments within the Lake Missoula basin (Chambers, 1971) that glacial Lake Mis-

soula drained repeatedly with at least 22 partial drainages of the lake to altitudes below 985 m. Chambers (1984), however, found "no evidence that each drainage was catastrophic." Depositional evidence of multiple water-level fluctuations is confined to only the lowest parts of the lake basin, thinning up the valley sides (Chambers, 1984). Chambers (1984) also noted that multiple cycles of lacustrine deposits are preserved within the troughs of giant current ripples in the Camas Prairie, further illustrating the disparity between a large, early release and later, lower-level fluctuations recorded by the cycles of lacustrine deposits. The volume of water available for flooding from a 1,080-m glacial Lake Missoula was about 1,055 km<sup>3</sup> (estimated from the hypsometric data of Clarke and others, 1984), less than 50% of the maximum lake volume. These observations are consistent with the view that only once in its history did the late Pleistocene Glacial Lake Missoula rise to the 1,265- to 1,280-m level and that the release from this level was more cataclysmic than subsequent releases.

**Conditions at the Ice Dam.** The failure mechanics of the ice dam depend on the local glacier profile, the geometry of the blockage, and the state of the glacier-bed interface. None of these conditions is known with certainty. The stratigraphic evidence for multiple, periodic, late Wisconsinan floods of systematically varying magnitude and frequency is consistent with a self-dumping, subglacial release mechanism for the flows that left these deposits. This mechanism has been observed for a number of historic jökulhlaups (for example, Thorarinsson, 1939; Post and Mayo, 1971; Sturm and others, 1987). Clarke and others (1984) and Waitt (1985), guided by these examples, quantitatively evaluated the possible conditions associated with such a failure mechanism for glacial Lake Missoula.

There are, however, mechanisms that could have resulted in a more rapid failure and a larger peak discharge. As Waitt (1985, p. 1282) noted, "preliminary calculations" (presumably associated with maximum lake-level conditions) indicate that the enlarging tunnel (due to dynamic thermal erosion) could expand to the point that the "tunnel roof collapsed and allowed most of glacial Lake Missoula to escape." Clarke and others (1984, p. 294) mentioned the possibility "that the tunnel roof collapses during the flood and the dam is swept away." The potential of such a failure is greater during a release from the maximum glacial Lake Missoula level because of the much greater amount of energy available for thermal erosion while the lake emptied

through the ice dam. The energy release associated with passage of the maximum (1,265-m stage) glacial Lake Missoula through the outlet is about three times that of a 1,080-m stage glacial Lake Missoula.

Another possibility is that the ice dam failed because the shear strength of the lobe of ice blocking the Clark Fork River was less than the hydrostatic force exerted by the impounded water. This hypothesis is difficult to evaluate because the geometry of a blocking lobe of ice is not known for the time of maximum lake level. At its maximum extent during late Wisconsinan time, the Purcell Trench lobe occupied the entire Lake Pend Oreille basin. With such an arrangement, the ice dam would probably have been stable against such a failure mechanism. At less advanced positions, however, the western margin of the glacier would likely have calved into the Pleistocene predecessor to Lake Pend Oreille. Under such conditions, the breadth of the blocking lobe of ice could have been reduced rapidly if it became ungrounded, leading to a brittle and rapid failure of the ice dam. This type of failure would be somewhat analogous to the largest known historic glacial outburst flood that occurred in October 1986, when Russell Lake burst through Hubbert Glacier in southeastern Alaska (Mayo, 1988, 1989).

## CONCLUSIONS

The field evidence for maximum Missoula Flood stages, in conjunction with step-back-water water-surface profile reconstructions, indicates a peak discharge of about 10 million m<sup>3</sup>·sec<sup>-1</sup> at Wallula Gap and a minimum of about 17 million m<sup>3</sup>·sec<sup>-1</sup> in the Spokane Valley (near the point of release). Considering uncertainties in the calculation parameters and the geologic evidence, the discharge at Wallula Gap can be limited to between 7.5 and 12 million m<sup>3</sup>·sec<sup>-1</sup>. The maximum discharge at the point of release is likely to have been larger than the 17 million m<sup>3</sup>·sec<sup>-1</sup> discharge estimated for the Spokane Valley.

The discharge estimate for the Spokane Valley is generally larger than independent analyses of the peak discharges of maximum glacial Lake Missoula outbursts derived from theoretical and empirical evaluations of a subglacial release mechanism. This leads us to propose that at least one failure of the ice dam may have been by a more cataclysmic mechanism than the subglacial jökulhlaups postulated (Waitt, 1980, 1984, 1985; Atwater, 1986) for emplacement of rhythmic strata at diverse sites throughout the

flooded region. The hypothesis of a larger, more cataclysmic, late Wisconsinan flood preceding multiple subglacial jökulhlaups is consistent with geologic evidence in the Channeled Scabland and in the glacial Lake Missoula basin.

Finally, it should be made clear that we are not attempting to disparage the well-constructed "case for periodic, colossal jökulhlaups." We are simply concerned that, in our view, anomalies still exist between some aspects of the field evidence and the conceptual models that have been advocated. The position that "the scores-of-floods hypothesis completes Bretz's imaginative theory" (Waitt, 1985, p. 1286) may prematurely divert attention from some of the outstanding problems that remain in interpreting the spectacular features of the Channeled Scabland.

## ACKNOWLEDGMENTS

This project was supported by the NASA Planetary Geology and Geophysics Program, Grant NAGW-285, and by the National Science Foundation Division of Earth Sciences, Surficial Processes Program, Grant EAR-8805321. Field assistance was provided by Karen A. Demsey. This paper has benefited from our discussions with Richard Waitt. G.S.A. reviewers Richard Waitt, Gary Clarke, Richard Craig, and Arthur Sylvester provided constructive critiques. Gary Smith, Sara Rathdrum, and Brian Atwater reviewed early versions of this manuscript.

## REFERENCES CITED

- Allison, I. S., 1933, New version of the Spokane flood: *Geological Society of America Bulletin*, v. 44, p. 675-722.
- Atwater, B. F., 1984, Periodic floods from glacial Lake Missoula into the Sanpoil arm of glacial Lake Columbia, northeastern Washington: *Geology*, v. 12, p. 464-467.
- , 1986, Pleistocene glacial-lake deposits of the Sanpoil River Valley, northeastern Washington: *U.S. Geological Survey Bulletin* 1661, 39 p.
- , 1987, Status of glacial Lake Columbia during the last floods from glacial Lake Missoula: *Quaternary Research*, v. 27, p. 182-201.
- Baker, V. R., 1973, Paleohydrology and sedimentology of Lake Missoula flooding in eastern Washington: *Geological Society of America Special Paper* 144, 79 p.
- Baker, V. R., and Bunker, R. C., 1985, Cataclysmic late Pleistocene flooding from glacial Lake Missoula: A review: *Quaternary Science Reviews*, v. 4, p. 1-41.
- Beget, J. E., 1986, Comment on "Outburst floods from glacial Lake Missoula" by G.K.C. Clarke, W. H. Mathews, and R. T. Pack: *Quaternary Research*, v. 25, no. 1, p. 136-138.
- Bjornstad, B. N., Focht, K. R., and Tallman, A. M., 1991, Quaternary stratigraphy of the Pasco Basin, south-central Washington, in Morrison, R. B., ed., *Quaternary nonglacial geology: Conterminous U.S.*: Boulder, Colorado, Geological Society of America, *The Geology of North America*, Volume K-2, p. 228-238.
- Bolke, E. L., and Vaccaro, J. J., 1981, Digital-model simulation of the hydrologic flow system, with emphasis on groundwater, in the Spokane Valley, Washington and Idaho: *U.S. Geological Survey Water Resources Investigations, Open-File Report* 80-1300, 43 p.
- Bretz, J. H., 1925, The Spokane flood beyond the Channeled Scablands: *Journal of Geology*, v. 33, p. 97-115, 236-259.
- , 1930, Valley deposits immediately west of the Channeled Scabland: *Journal of Geology*, v. 38, p. 385-422.
- , 1969, The Lake Missoula floods and the Channeled Scabland: *Journal of Geology*, v. 77, p. 505-543.

- Bunker, R. C., 1980, Catastrophic flooding in the Badger Coulee area, south-central Washington [M.A. thesis] Austin, Texas, University of Texas, 184 p.
- Chambers, R. L., 1971, Sedimentation in glacial Lake Missoula [M.S. thesis] Missoula, Montana, University of Montana, 100 p.
- , 1984, Sedimentary evidence for multiple glacial Lakes Missoula: Montana Geological Field Conference on Northwestern Montana, Kalispell, Montana, p. 189-199.
- Chow, V. T., 1959, *Open-channel hydraulics*: New York, McGraw-Hill, 690 p.
- Clarke, G.K.C., 1982, Glacier outburst floods from "Hazard Lake," Yukon Territory, and the problem of flood magnitude prediction: *Journal of Glaciology*, v. 28, p. 3-21.
- Clarke, G.K.C., Mathews, W. H., and Pack, R. T., 1984, Outburst floods from glacial Lake Missoula: *Quaternary Research*, v. 22, p. 289-299.
- Costa, J. E., 1988, Floods from dam failures, in Baker, V. R., Koebel, R. C., and Patton, P. C., eds., *Flood geomorphology*: New York, John Wiley and Sons, p. 439-463.
- Craig, R. G., 1987, Dynamics of a Missoula flood, in Mayer, L., and Nash, D., eds., *Catastrophic flooding*: Boston, Massachusetts, Allen and Unwin, p. 305-332.
- Craig, R. G., and Hanson, J. P., 1985, Erosion potential from Missoula floods in the Pasco Basin, Washington: Battelle Pacific Northwest Laboratory Document PNL-5684, UC-70, 185 p. and 5 appendices.
- Dalrymple, T., and Benson, M. A., 1967, Measurement of peak discharge by the slope-area method: U.S. Geological Survey Techniques Water-Resources Investigations, Book 3, Chap. A2, 12 p.
- Ely, L. L., and Baker, V. R., 1985, Reconstructing paleoflood hydrology with slackwater deposits, Verde River, Arizona: *Physical Geography*, v. 5, p. 103-126.
- Feldman, A. D., 1981, HEC models for water resources system simulation, theory, and experience: *Advances in Hydroscience*, v. 12, p. 297-423.
- Flint, R. F., 1938, Origin of the Cheney-Palouse scabland tract: *Geological Society of America Bulletin*, v. 49, p. 461-524.
- Hanson, L. G., 1970, The origin and deformation of Moses Coulee and other scabland features on the Waterville Plateau, Washington [Ph.D. thesis]: Seattle, Washington, University of Washington, 137 p.
- Hydrologic Engineering Center, 1985, HEC-2 water surface profiles users manual: Davis, California, The Hydrologic Engineering Center (U.S. Army Corps of Engineers), 37 p.
- Jarrett, R. D., and Malde, H. E., 1987, Paleodischarge of the late Pleistocene Bonneville Flood, Snake River, Idaho, computed from new evidence: *Geological Society of America Bulletin*, v. 99, p. 127-134.
- Kiver, E. P., and Stradling, D. F., 1982, Quaternary geology of the Spokane area, in Roberts, S., and Fountain, D., eds., 1980 Field conference guidebook: Spokane, Washington, Tobacco Root Geological Society, p. 26-44.
- , 1989, The Spokane Valley and northern Columbia Plateau, in Breckenridge, R. M., ed., *Glacial Lake Missoula and the Channeled Scabland*: International Geological Congress, 28th, Field Trip Guidebook T310, p. 23-36.
- Mayo, L. R., 1988, Advance of Hubbard Glacier and closure of Russell Fjord, Alaska—Environmental effects and hazards in Yakutat area, in Galloway, J. P., and Hamilton, T. D., eds., *Geologic studies in Alaska by the U.S. Geological Survey during 1987*: U.S. Geological Survey Circular 1016, p. 4-16.
- , 1989, Advance of Hubbard Glacier and 1986 outburst of Russell Fjord, Alaska, U.S.A.: *Annals of Glaciology*, v. 13, p. 189-194.
- Newcomb, R. C., 1958, Ringold Formation of Pleistocene age in type locality, the White Bluffs, Washington: *American Journal of Science*, v. 256, p. 328-340.
- , 1969, Effect of tectonic structure on the occurrence of ground water in the basalt of the Columbia River Group of The Dalles area, Oregon and Washington: U.S. Geological Survey Professional Paper 383-C, 33 p.
- Newcomb, R. C., Strand, J. R., and Frank, F. J., 1972, Geology and ground-water characteristics of the Hanford Reservation of the U.S. Atomic Energy Commission, Washington: U.S. Geological Survey Professional Paper 717, 78 p.
- O'Connor, J. E., 1990, Hydrology, hydraulics, and sediment transport of Pleistocene Lake Bonneville flooding on the Snake River, Idaho [Ph.D. thesis]: Tucson, Arizona, University of Arizona, 192 p.
- O'Connor, J. E., and Webb, R. H., 1988, Hydraulic modeling for paleoflood analysis, in Baker, V. R., Koebel, R. C., and Patton, P. C., eds., *Flood geomorphology*: New York, John Wiley and Sons, p. 403-418.
- Pardee, J. T., 1942, Unusual currents in glacial Lake Missoula: *Geological Society of America Bulletin*, v. 53, p. 1569-1600.
- Post, A., and Mayo, L. R., 1971, Glacier-dammed lakes and outburst floods in Alaska: U.S. Geological Survey Hydrologic Investigations Atlas HA-455.
- Richmond, G. M., Fryxell, R., Neff, G. E., and Weis, P. L., 1965, The Cordilleran Ice Sheet of the northern Rocky Mountains and related Quaternary history of the Columbia Plateau, in Wright, H. E., Jr., and Frey, D. G., eds., *The Quaternary of the United States*: Princeton, New Jersey, Princeton University Press, p. 231-242.
- Sturm, M., Beget, J., and Benson, C., 1987, Observations of jökulhlaups from ice-dammed Strandine Lake, Alaska: Implications for paleohydrology, in Mayer, L., and Nash, D., eds., *Catastrophic flooding*: Boston, Massachusetts, Allen and Unwin, p. 79-94.
- Thorarinsson, S., 1939, The ice-dammed lakes of Iceland with particular reference to their value as indicators of glacier oscillations: *Geografiska Annaler*, v. 21, p. 216-242.
- Wait, R. B., 1977, Missoula flood sizes Okanogan lobe: *Geological Society of America Abstracts with Programs*, v. 9, p. 770.
- , 1980, About forty last-glacial Lake Missoula jökulhlaups through southern Washington: *Journal of Geology*, v. 88, p. 653-679.
- , 1982, Surficial geology, in Tabor, R. W., and others, *Geologic map of the Wenatchee 1:100,000 quadrangle, Washington*: U.S. Geological Survey Miscellaneous Investigations Map I-1311.
- , 1984, Periodic jökulhlaups from Pleistocene glacial Lake Missoula: New evidence from varved sediment in northern Idaho and Washington: *Quaternary Research*, v. 22, p. 46-58.
- , 1985, Case for periodic, colossal jökulhlaups from Pleistocene glacial Lake Missoula: *Geological Society of America Bulletin*, v. 96, p. 1271-1286.
- , 1987, Erosional landscapes and surficial deposits, in Tabor, R. W., Frizzell, V. A., Jr., Whetten, J. T., Wait, R. B., Swanson, D. A., Byerly, G. R., Booth, D. B., Hetherington, M. J., and Zartman, R. E., *Geologic map of the Chelan 30-minute by 60-minute quadrangle, Washington*: U.S. Geological Survey Miscellaneous Investigations Map I-1661.
- Wait, R. B., and Swanson, D. A., 1987, Geomorphic evolution of the Columbia Plain and River, in Graf, W. L., ed., *Geomorphic systems of North America*: Boulder, Colorado, Geological Society of America Centennial Special Volume 2, p. 403-416.
- Wait, R. B., and Thorson, R. M., 1983, The Cordilleran Ice Sheet in Washington, Idaho, and Montana, in Wright, H. E., Jr., ed., *Late Quaternary environments of the United States; Volume I, The late Pleistocene*: Minneapolis, Minnesota, University of Minnesota Press, p. 53-70.

MANUSCRIPT RECEIVED BY THE SOCIETY OCTOBER 22, 1990  
 REVISED MANUSCRIPT RECEIVED JULY 10, 1991  
 MANUSCRIPT ACCEPTED JULY 22, 1991